



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12P 21/06, C12N 9/02, 1/20, 15/00, C07H 21/04, C07K 1/00	A1	(11) International Publication Number: WO 99/23242 (43) International Publication Date: 14 May 1999 (14.05.99)
(21) International Application Number: PCT/US98/23419 (22) International Filing Date: 3 November 1998 (03.11.98) (30) Priority Data: 60/064,195 4 November 1997 (04.11.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indi- anapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): COSTELLO, Colleen, An- gela [US/US]; 733 Lake Drive, Lawrenceville, NJ 08648 (US). HERSHBERGER, Charles, Lee [US/US]; 1475 Har- mony Trail West, Greenfield, IN 46140 (US). MENKE, Michael, Andrew [US/US]; 345 North Lesley Avenue, Indi- anapolis, IN 46219 (US). ZMIJEWSKI, Milton, Joseph, Jr. [US/US]; 10152 Partridge Place, Carmel, IN 46033 (US). (74) Agents: WEBSTER, Thomas, D. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: KETOREDUCTASE GENE AND PROTEIN FROM YEAST (57) Abstract This invention provides a cloned ketoreductase gene, vectors for expressing same, recombinant host cells that express said vector-borne gene, and a method for stereospecifically reducing a ketone using a recombinant ketoreductase, or a recombinant host cell that expresses a cloned ketoreductase gene.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

KETOREDUCTASE GENE AND PROTEIN FROM YEAST

CROSS-REFERENCE

5 This application claims the benefit of U.S. Provisional Application No. 60/064,195, filed November 4, 1997.


FIELD OF THE INVENTION

10 This invention relates to recombinant DNA technology. In particular the invention pertains to the cloning of a ketoreductase gene from *Zygosaccharomyces rouxii*, and the use of recombinant hosts expressing fungal ketoreductase genes in a process for stereospecific reduction of ketones.

BACKGROUND OF THE INVENTION

15 2,3 Benzodiazepine derivatives are potent antagonists of the AMPA (α -amino-3-hydroxy-5 methylisoxazole-4-propionic acid) class of receptors in the
20 mammalian central nervous system (See I. Tarnawa et al. In *Amino Acids: Chemistry, Biology and Medicine*, Eds. Lubec and Rosenthal, Leiden, 1990). These derivative compounds have potentially widespread applications as neuroprotective agents, particularly as anti-convulsants. One series of 2,3
25 benzodiazepines is considered particularly advantageous for such use, and this series of compounds has the following general formula:

-2-



- 5 Wherein R is hydrogen or C₁-C₁₀ alkyl; and
 X is hydrogen, C₁-C₁₀ alkyl, acyl, aryl, amido or
carboxyl, or a substituted derivative thereof.

The clinical potential for these compounds has led to
interest in developing more efficient synthetic methods.

- 10 Biologically-based methods in which a ketoreductase enzyme
provides a stereospecific reduction in a whole-cell process
using fungal cells have been described in U.S. Patent
application serial number 08/413,036.

15 BRIEF SUMMARY OF THE INVENTION

- The present invention provides isolated nucleic acid
molecules that encode a ketoreductase enzyme from *Z. rouxii*..
The invention also provides the protein product of said
nucleic acid, in substantially purified form. Also provided
20 are methods for the formation of chiral alcohols using a
purified ketoreductase enzyme, or a recombinant host cell
that expresses a fungal ketoreductase gene.

- Having the cloned ketoreductase gene enables the
production of recombinant ketoreductase protein, and the
25 production of recombinant host cells expressing said

-3-

protein, wherein said recombinant cells can be used in a stereospecific reduction of ketones.

In one embodiment the present invention relates to an isolated DNA molecule encoding ketoreductase protein, said
5 DNA molecule comprising the nucleotide sequence identified as SEQ ID NO:1.

In another embodiment the present invention relates to a substantially purified ketoreductase protein molecule from *Z. rouxii*.

10 In another embodiment the present invention relates to a ketoreductase protein molecule from *Z. rouxii*, wherein said protein molecule comprises the sequence identified as SEQ ID NO:2.

In a further embodiment the present invention relates
15 to a ribonucleic acid molecule encoding ketoreductase protein, said ribonucleic acid molecule comprising the sequence identified as SEQ ID NO:3.

In yet another embodiment, the present invention relates to a recombinant DNA vector that incorporates a
20 ketoreductase gene in operable-linkage to gene expression sequences, enabling said gene to be transcribed and translated in a host cell.

In still another embodiment the present invention relates to host cells that have been transformed or
25 transfected with a cloned ketoreductase gene such that said ketoreductase gene is expressed in the host cell.

In a still further embodiment, the present invention relates to a method for producing chiral alcohols using recombinant host cells that express an exogenously
30 introduced ketoreductase gene.

-4-

In yet another embodiment, the present invention relates to a method for producing chiral alcohols using recombinant host cells that have been transformed or transfected with a ketoreductase gene from *Z. rouxii*, or *S. cerevisiae*.

In yet another embodiment, the present invention relates to a method for producing chiral alcohols using a purified fungal ketoreductase.

10

DETAILED DESCRIPTION OF THE INVENTION

Definitions

SEQ ID NO:1 - SEQ ID NO:3 comprises the DNA, protein, and RNA sequences of ketoreductase from *Z. rouxii*.

15

SEQ ID NO:4- SEQ ID NO:6 comprises the DNA, protein, and RNA sequences of gene YDR541c from *S. cerevisiae*.

SEQ ID NO:7- SEQ ID NO:9 comprises the DNA, protein, and RNA sequences of YOL151w from *S. cerevisiae*.

20

SEQ ID NO:10- SEQ ID NO:12 comprises the DNA, protein, and RNA sequences of YGL157w from *S. cerevisiae*.

SEQ ID NO:13- SEQ ID NO:15 comprises the DNA, protein, and RNA sequences of YGL039w from *S. cerevisiae*.

25

The term "fusion protein" denotes a hybrid protein molecule not found in nature comprising a translational fusion or enzymatic fusion in which two or more different proteins or fragments thereof are covalently linked on a single polypeptide chain.

30

The term "plasmid" refers to an extrachromosomal genetic element. The starting plasmids herein are either commercially available, publicly available on an

-5-

unrestricted basis, or can be constructed from available plasmids in accordance with published procedures. In addition, equivalent plasmids to those described are known in the art and will be apparent to the ordinarily skilled
5 artisan.

"Recombinant DNA cloning vector" as used herein refers to any autonomously replicating agent, including, but not limited to, plasmids and phages, comprising a DNA molecule to which one or more additional DNA segments can or
10 have been added.

The term "recombinant DNA expression vector" or "expression vector" as used herein refers to any recombinant DNA cloning vector, for example a plasmid or phage, in which a promoter and other regulatory elements are present thereby
15 enabling transcription of an inserted DNA.

The term "vector" as used herein refers to a nucleic acid compound used for introducing exogenous DNA into host cells. A vector comprises a nucleotide sequence which may encode one or more protein molecules. Plasmids,
20 cosmids, viruses, and bacteriophages, in the natural state or which have undergone recombinant engineering, are examples of commonly used vectors.

The terms "complementary" or "complementarity" as used herein refers to the capacity of purine and pyrimidine
25 nucleotides to associate through hydrogen bonding in double stranded nucleic acid molecules. The following base pairs are complementary: guanine and cytosine; adenine and thymine; and adenine and uracil. As used herein

"complementary" means that at least one of two hybridizing
30 strands is fully base-paired with the other member of said

-6-

hybridizing strands, and there are no mismatches. Moreover, at each nucleotide position of said one strand, an "A" is paired with a "T", a "T" is paired with an "A", a "G" is paired with a "C", and a "C" is paired with a "G".

5 "Isolated nucleic acid compound" refers to any RNA or DNA sequence, however constructed or synthesized, which is locationally distinct from its natural location.

A "primer" is a nucleic acid fragment which functions as an initiating substrate for enzymatic or
10 synthetic elongation of, for example, a nucleic acid molecule.

The term "promoter" refers to a DNA sequence which directs transcription of DNA to RNA. An inducible promoter is one that is regulatable by environmental signals, such as
15 carbon source, heat, metal ions, chemical inducers, etc.; a constitutive promoter generally is expressed at a constant level and is not regulatable.

A "probe" as used herein is a labeled nucleic acid compound which can hybridize with another nucleic acid
20 compound.

The term "hybridization" as used herein refers to a process in which a single-stranded nucleic acid molecule joins with a complementary strand through nucleotide base pairing. "Selective hybridization" refers to hybridization
25 under conditions of high stringency. The degree of hybridization depends upon, for example, the degree of complementarity, the stringency of hybridization, and the length of hybridizing strands.

"Substantially identical" means a sequence having
30 sufficient homology to hybridize under stringent conditions

-7-

and/or be at least 90% identical to a sequence disclosed herein.

The term "stringency" relates to nucleic acid hybridization conditions. High stringency conditions disfavor non-homologous base pairing. Low stringency conditions have the opposite effect. Stringency may be altered, for example, by changes in temperature, denaturants, and salt concentration. Typical high stringency conditions comprise hybridizing at 50°C to 65°C in 5X SSPE and 50% formamide, and washing at 50°C to 65°C in 0.5X SSPE; typical low stringency conditions comprise hybridizing at 35°C to 37° in 5X SSPE and 40% to 45% formamide and washing at 42°C in 1X-2X SSPE.

"SSPE" denotes a hybridization and wash solution comprising sodium chloride, sodium phosphate, and EDTA, at pH 7.4. A 20X solution of SSPE is made by dissolving 174 g of NaCl, 27.6 g of $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$, and 7.4 g of EDTA in 800 ml of H_2O . The pH is adjusted with NaOH and the volume brought to 1 liter.

"SSC" denotes a hybridization and wash solution comprising sodium chloride and sodium citrate at pH 7. A 20X solution of SSC is made by dissolving 175 g of NaCl and 88 g of sodium citrate in 800 ml of H_2O . The volume is brought to 1 liter after adjusting the pH with 10N NaOH.

The ketoreductase gene encodes a novel enzyme that catalyzes an asymmetric reduction of selected ketone substrates (See Equation 1 and Table 1).

Equation 1

- 8 -















Table 1: Substrate specificity of ketoreductase from *Z. rouxii*.

Compound	Concentration (mM)	% Relative Activity	Compound	Concentration (mM)	% Relative Activity
 1	3	100	 9	3	194
 4	5	18	 10	0.8	86
 5	5	42	 11	0.6	17
 6	4	37	 12	5	100
 7	0.6	4	 13	5	32
 8	0.6	0			

The ketoreductase enzymes disclosed herein are members of the carbonyl reductase enzyme class. Carbonyl reductases are involved in the reduction of xenobiotic carbonyl compounds (Hara et. al, Arch. Biochem. Biophys., 244, 238-

-10-

247, 1986) and have been classified into the short-chain dehydrogenase/reductase (SDR) enzyme superfamily (Jörnvall et. al, *Biochemistry*, 34, 6003-6013, 1995) and the single-domain reductase/epimerase/dehydrogenase (RED) enzyme superfamily (Labesse et. al, *Biochem. J.*, 304, 95-99, 1994). The ketoreductases of this invention are able to effectively reduce a variety of α -ketolactones, α -ketolactams, and diketones (Table 1).

The ketoreductase gene of *Z. rouxii* comprises a DNA sequence designated herein as SEQ ID NO:1. Those skilled in the art will recognize that owing to the degeneracy of the genetic code (i.e. 64 codons which encode 20 amino acids), numerous "silent" substitutions of nucleotide base pairs could be introduced into the sequence identified as SEQ ID NO:1 without altering the identity of the encoded amino acid(s) or protein product. All such substitutions are intended to be within the scope of the invention.

Gene Isolation Procedures

Those skilled in the art will recognize that the ketoreductase gene may be obtained by a plurality of applicable recombinant DNA techniques including, for example, polymerase chain reaction (PCR) amplification, hybridization to a genomic or cDNA library, or de novo DNA synthesis. (See e.g., J.Sambrook et al. Molecular Cloning, 2d Ed. Chap. 14 (1989)).

Methods for constructing cDNA libraries in a suitable vector such as a plasmid or phage for propagation in procaryotic or eucaryotic cells are well known to those

-11-

skilled in the art. [See e.g. J.Sambrook et al. Supra].
Suitable cloning vectors are widely available.

Skilled artisans will recognize that the ketoreductase gene or fragment thereof could be isolated by PCR amplification from a human cDNA library prepared from a tissue in which said gene is expressed, using oligonucleotide primers targeted to any suitable region of SEQ ID NO:1. Methods for PCR amplification are widely known in the art. See e.g. PCR Protocols: A Guide to Method and Application, Ed. M. Innis et.al., Academic Press (1990). The amplification reaction comprises template DNA, suitable enzymes, primers, nucleoside triphosphates, and buffers, and is conveniently carried out in a DNA Thermal Cycler (Perkin Elmer Cetus, Norwalk, CT). A positive result is determined by detecting an appropriately-sized DNA fragment following gel electrophoresis.

Protein Production Methods

One embodiment of the present invention relates to the substantially purified ketoreductase enzyme (identified herein as SEQ ID NO:2) encoded by the *Z. rouxii* ketoreductase gene (identified herein as SEQ ID NO:1).

Skilled artisans will recognize that the proteins of the present invention can be synthesized by a number of different methods, such as chemical methods well known in the art, including solid phase peptide synthesis or recombinant methods. Both methods are described in U.S. Patent 4,617,149, incorporated herein by reference. The proteins of the invention can also be purified by well known

-12-

methods from a culture of cells that produce the protein, for example, *Z. rouxii*.

The principles of solid phase chemical synthesis of polypeptides are well known in the art and may be found in general texts in the area. See, e.g., H. Dugas and C. Penney, Bioorganic Chemistry (1981) Springer-Verlag, New York, 54-92. For example, peptides may be synthesized by solid-phase methodology utilizing an Applied Biosystems 430A peptide synthesizer (Applied Biosystems, Foster City, CA) and synthesis cycles supplied by Applied Biosystems.

The protein of the present invention can also be produced by recombinant DNA methods using the cloned ketoreductase gene. Recombinant methods are preferred if a high yield is desired. Expression of the cloned gene can be carried out in a variety of suitable host cells, well known to those skilled in the art. For this purpose, the ketoreductase gene is introduced into a host cell by any suitable means, well known to those skilled in the art. While chromosomal integration of the cloned gene is within the scope of the present invention, it is preferred that the gene be cloned into a suitable extra-chromosomally maintained expression vector so that the coding region of the ketoreductase gene is operably-linked to a constitutive or inducible promoter.

The basic steps in the recombinant production of the ketoreductase protein are:

- a) constructing a natural, synthetic or semi-synthetic DNA encoding ketoreductase protein;

-13-

b) integrating said DNA into an expression vector in a manner suitable for expressing the ketoreductase protein, either alone or as a fusion protein; or integrating said DNA into a host chromosome such that said DNA expresses ketoreductase;

c) transforming or otherwise introducing said vector into an appropriate eucaryotic or prokaryotic host cell forming a recombinant host cell,

d) culturing said recombinant host cell in a manner to express the ketoreductase protein; and

e) recovering and substantially purifying the ketoreductase protein by any suitable means, well known to those skilled in the art.

Expressing Recombinant ketoreductase Protein in Procaryotic and Eucaryotic Host Cells

Procaryotes may be employed in the production of the ketoreductase protein. For example, the *Escherichia coli* K12 strain 294 (ATCC No. 31446) or strain RV308 is particularly useful for the prokaryotic expression of foreign proteins. Other strains of *E. coli*, bacilli such as *Bacillus subtilis*, enterobacteriaceae such as *Salmonella typhimurium* or *Serratia marcescans*, various *Pseudomonas*

-14-

species and other bacteria, such as *Streptomyces*, may also be employed as host cells in the cloning and expression of the recombinant proteins of this invention.

Promoter sequences suitable for driving the expression of genes in procaryotes include β -lactamase [e.g. vector pGX2907, ATCC 39344, contains a replicon and β -lactamase gene], lactose systems [Chang et al., *Nature* (London), 275:615 (1978); Goeddel et al., *Nature* (London), 281:544 (1979)], alkaline phosphatase, and the tryptophan (trp) promoter system [vector PATH1 (ATCC 37695) which is designed to facilitate expression of an open reading frame as a trpE fusion protein under the control of the trp promoter]. Hybrid promoters such as the tac promoter (isolatable from plasmid pDR540, ATCC-37282) are also suitable. Still other bacterial promoters, whose nucleotide sequences are generally known, enable one of skill in the art to ligate such promoter sequences to DNA encoding the proteins of the instant invention using linkers or adapters to supply any required restriction sites. Promoters for use in bacterial systems also will contain a Shine-Dalgarno sequence operably linked to the DNA encoding the desired polypeptides. These examples are illustrative rather than limiting.

The protein(s) of this invention may be synthesized either by direct expression or as a fusion protein comprising the protein of interest as a translational fusion with another protein or peptide which may be removable by enzymatic or chemical cleavage. It is often observed in the production of certain peptides in recombinant systems that expression as a fusion protein

-15-

prolongs the lifespan, increases the yield of the desired peptide, or provides a convenient means of purifying the protein. A variety of peptidases (e.g. enterokinase and thrombin) which cleave a polypeptide at specific sites or
5 digest the peptides from the amino or carboxy termini (e.g. diaminopeptidase) of the peptide chain are known. Furthermore, particular chemicals (e.g. cyanogen bromide) will cleave a polypeptide chain at specific sites. The skilled artisan will appreciate the modifications necessary
10 to the amino acid sequence (and synthetic or semi-synthetic coding sequence if recombinant means are employed) to incorporate site-specific internal cleavage sites. See e.g., P. Carter, "Site Specific Proteolysis of Fusion
15 Molecular Mechanisms to Large Scale Processes, American Chemical Society, Washington, D.C. (1990).

In addition to procaryotes, a variety of eucaryotic microorganisms including yeast are suitable host cells. The yeast *Saccharomyces cerevisiae* is the most
20 commonly used eucaryotic microorganism. Other yeasts such as *Kluyveromyces lactis*, *Schizosaccharomyces pombe*, and *Pichia pastoris* are also suitable. For expression in *Saccharomyces*, the plasmid YRp7 (ATCC-40053), for example, may be used. See, e.g., L. Stinchcomb, et al., *Nature*,
25 282:39 (1979); J. Kingsman et al., *Gene*, 7:141 (1979); S. Tschemper et al., *Gene*, 10:157 (1980). Plasmid YRp7 contains the TRP1 gene which provides a selectable marker for use in a *trp1* auxotrophic mutant.

30 Purification of Recombinantly-Produced ketoreductase Protein

-16-

An expression vector carrying a cloned ketoreductase gene is transformed or transfected into a suitable host cell using standard methods. Host cells may comprise procaryotes, such as *E. coli*, or simple eucaryotes, such as *Z. rouxii*, *S. cerevisiae*, *S. pombe*, *P. pastoris*, and *K. Lactis*. Cells which contain the vector are propagated under conditions suitable for expression of an encoded ketoreductase protein. If the recombinant gene has been placed under the control of an inducible promoter then suitable growth conditions would incorporate the appropriate inducer. The recombinantly-produced protein may be purified from cellular extracts of transformed cells by any suitable means.

In a preferred process for protein purification, the ketoreductase gene is modified at the 5' end to incorporate several histidine residues at the amino terminus of the ketoreductase protein product. This "histidine tag" enables a single-step protein purification method referred to as "immobilized metal ion affinity chromatography" (IMAC), essentially as described in U.S. Patent 4,569,794 which hereby is incorporated by reference. The IMAC method enables rapid isolation of substantially pure ketoreductase protein starting from a crude cellular extract.

Other embodiments of the present invention comprise isolated nucleic acid sequences which encode SEQ ID NO:2. As skilled artisans will recognize, the amino acid compounds of the invention can be encoded by a multitude of different nucleic acid sequences because most of the amino acids are encoded by more than one codon. Because these alternative nucleic acid sequences would encode the same

-17-

amino acid sequences, the present invention further comprises these alternate nucleic acid sequences.

The ketoreductase genes disclosed herein, for example SEQ ID NO:1, may be produced using synthetic methodology. The synthesis of nucleic acids is well known in the art. See, e.g., E.L. Brown, R. Belagaje, M.J. Ryan, and H.G. Khorana, Methods in Enzymology, 68:109-151 (1979). A DNA segment corresponding to a ketoreductase gene could be generated using a conventional DNA synthesizing apparatus, such as the Applied Biosystems Model 380A or 380B DNA synthesizers (Applied Biosystems, Inc., 850 Lincoln Center Drive, Foster City, CA 94404) which employ phosphoramidite chemistry. Alternatively, phosphotriester chemistry may be employed to synthesize the nucleic acids of this invention. [See, e.g., M.J. Gait, ed., Oligonucleotide Synthesis, A Practical Approach, (1984).]

In an alternative methodology, namely PCR, a DNA sequence comprising a portion or all of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, or SEQ ID NO:13 can be generated from a suitable DNA source, for example *Z. rouxii* or *S. cerevisiae* genomic DNA or cDNA. For this purpose, suitable oligonucleotide primers targeting SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, SEQ ID NO:13 or region therein are prepared, as described in U.S. Patent No. 4,889,818, which hereby is incorporated by reference. Protocols for performing the PCR are disclosed in, for example, PCR Protocols: A Guide to Method and Applications, Ed. Michael A. Innis et al., Academic Press, Inc. (1990).

The ribonucleic acids of the present invention may be prepared using the polynucleotide synthetic methods

-18-

discussed *supra*, or they may be prepared enzymatically using RNA polymerase to transcribe a ketoreductase DNA template. See e.g., J. Sambrook, et. al., *supra*, at 18.82-18.84.

This invention also provides nucleic acids, RNA or
5 DNA, which are complementary to SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, or SEQ ID NO:15.

The present invention also provides probes and primers useful for a variety of molecular biology techniques
10 including, for example, hybridization screens of genomic, subgenomic, or cDNA libraries. A nucleic acid compound comprising SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, or SEQ ID NO:15, or a complementary sequence
15 thereof, or a fragment thereof, which is at least 18 base pairs in length, and which will selectively hybridize to DNA encoding a ketoreductase, is provided. Preferably, the 18 or more base pair compound is DNA. See e.g. B. Wallace and G. Miyada, "Oligonucleotide Probes for the Screening of
20 Recombinant DNA Libraries," In Methods in Enzymology, Vol. 152, 432-442, Academic Press (1987).

Probes and primers can be prepared by enzymatic methods well known to those skilled in the art (See e.g. Sambrook et al. *supra*). In a most preferred embodiment
25 these probes and primers are synthesized using chemical means as described above.

Another aspect of the present invention relates to recombinant DNA cloning vectors and expression vectors comprising the nucleic acids of the present invention. The
30 preferred nucleic acid vectors are those which comprise DNA.

-19-

The most preferred recombinant DNA vectors comprise a isolated DNA sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, or SEQ ID NO:13.

5 The skilled artisan understands that choosing the most appropriate cloning vector or expression vector depends upon a number of factors including the availability of restriction enzyme sites, the type of host cell into which the vector is to be transfected or transformed, the purpose
10 of the transfection or transformation (e.g., stable transformation as an extrachromosomal element, or integration into the host chromosome), the presence or absence of readily assayable or selectable markers (e.g., antibiotic resistance and metabolic markers of one type and
15 another), and the number of copies of the gene to be present in the host cell.

Vectors suitable to carry the nucleic acids of the present invention comprise RNA viruses, DNA viruses, lytic bacteriophages, lysogenic bacteriophages, stable
20 bacteriophages, plasmids, viroids, and the like. The most preferred vectors are plasmids.

When preparing an expression vector the skilled artisan understands that there are many variables to be considered, for example, whether to use a constitutive or
25 inducible promoter. Inducible promoters are preferred because they enable high level, regulatable expression of an operably-linked gene. Constitutive promoters are further suitable in instances for which secretion or extra-cellular export is desirable. The skilled artisan will recognize a
30 number of inducible promoters which respond to a variety of

-20-

inducers, for example, carbon source, metal ions, and heat. The practitioner also understands that the amount of nucleic acid or protein to be produced dictates, in part, the selection of the expression system. The addition of certain
5 nucleotide sequences is useful for directing the localization of a recombinant protein. For example, a sequence encoding a signal peptide preceding the coding region of a gene, is useful for directing the extra-cellular export of a resulting polypeptide.

10 Host cells harboring the nucleic acids disclosed herein are also provided by the present invention. Suitable host cells include procaryotes, such as *E. coli*, or simple eucaryotes, such as fungal cells, which have been transfected or transformed with a vector which comprises a
15 nucleic acid of the present invention.

The present invention also provides a method for constructing a recombinant host cell capable of expressing SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, or SEQ ID NO:14, said method comprising transforming or otherwise
20 introducing into a host cell a recombinant DNA vector that comprises an isolated DNA sequence which encodes SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, or SEQ ID NO:14. Preferred vectors for expression are those which comprise SEQ ID NO:1. Transformed host cells may be cultured
25 under conditions well known to skilled artisans such that SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, or SEQ ID NO:14 is expressed, thereby producing a ketoreductase protein in the recombinant host cell.

For the purpose of identifying or developing
30 inhibitors or other modifiers of the enzymes disclosed

-21-

herein, or for identifying suitable substrates for bioconversion, it would be desirable to identify compounds that bind and/or inhibit, or otherwise modify, the ketoreductase enzyme and its associated activity. A method
5 for determining agents that will modify the ketoreductase activity comprises contacting the ketoreductase protein with a test compound and monitoring the alteration of enzyme activity by any suitable means.

The instant invention provides such a screening
10 system useful for discovering compounds which bind the ketoreductase protein, said screening system comprising the steps of:

- a) preparing ketoreductase protein;
- b) exposing said ketoreductase protein to a test
15 compound;
- c) quantifying a modulation of activity by said compound.

Utilization of the screening system described above provides a means to determine compounds which may
20 alter the activity of ketoreductase. This screening method may be adapted to automated procedures such as a PANDEX[®] (Baxter-Dade Diagnostics) system, allowing for efficient high-volume screening of potential modifying agents.

In such a screening protocol, ketoreductase is
25 prepared as described herein, preferably using recombinant DNA technology. A test compound is introduced into a reaction vessel containing ketoreductase, followed by addition of enzyme substrate. For convenience the reaction can be coupled to the oxidation of NADPH, thereby enabling

-22-

progress to be monitored spectrophotometrically by measuring the absorbance at 340 nm. Alternatively, substrate may be added simultaneously with a test compound. In one method radioactively or chemically-labeled compound may be used.

- 5 The products of the enzymatic reaction are assayed for the chemical label or radioactivity by any suitable means. The absence or diminution of the chemical label or radioactivity indicates the degree to which the reaction is inhibited.

The following examples more fully describe the present invention. Those skilled in the art will recognize that the particular reagents, equipment, and procedures described are merely illustrative and are not intended to limit the present invention in any manner.

15

EXAMPLE 1

Construction of a DNA Vector for Expressing a Ketoreductase Gene in a Homologous or Heterologous Host

A plasmid comprising the *Z. rouxii* ketoreductase gene suitable for expressing said gene in a host cell, for example *E. coli* (DE3) strains, contains an origin of replication (Ori), an ampicillin resistance gene (Amp), useful for selecting cells which have incorporated the vector following a transformation procedure, and further comprises the *lacI* gene for repression of the *lac* operon, as well as the T7 promoter and T7 terminator sequences in operable linkage to the coding region of the ketoreductase gene. Parent plasmid pET11A (obtained from Novogen, Madison, WI) was linearized by digestion with endonucleases *NdeI* and *BamHI*. Linearized pET11A was ligated to a DNA fragment

-23-

bearing *NdeI* and *BamHI* sticky ends and further comprising the coding region of the *Z. rouxii* ketoreductase gene.

The ketoreductase gene is isolated most conveniently by the PCR. Genomic DNA from *Z. rouxii* isolated by standard methods was used for amplification of the ketoreductase gene. Primers are synthesized corresponding to the 5' and 3' ends of the gene (SEQ ID NO:1) to enable amplification of the coding region.

The ketoreductase gene (nucleotides 164 through 1177 of SEQ ID NO:1) ligated into the vector was modified at the 5' end (amino terminus of encoded protein) in order to simplify purification of the encoded ketoreductase protein. For this purpose, an oligonucleotide encoding 8 histidine residues and a factor Xa cleavage site was inserted after the ATG start codon at nucleotide positions 164 to 166 of SEQ ID NO:1. Placement of the histidine residues at the amino terminus of the encoded protein does not affect its activity and serves only to enable the IMAC one-step protein purification procedure.

20

EXAMPLE 2

Purification of Ketoreductase from *Z. rouxii*

Approximately 1 gram of *Z. rouxii* cell paste was resuspended in Lysing Buffer, comprising 50 mM Tris-Cl pH 7.5, 2 mM EDTA supplemented with pepstatin (1 µg/mL), leupeptin (1.25 µg/mL), aprotinin (2.5 µg/mL), and AEBSF (25 µg/mL). The cells were lysed using a DynoMill (GlenMills, Inc. Clifton, NJ) equipped with 0.5-0.75 mm lead free beads under continuous flow conditions according to the manufacturer's recommended use. After four complete passes

30

-24-

through the DynoMill, the material was centrifuged twice (25,000 x g for 30 minutes at 4°C). Solid ammonium sulfate (291 g/liter) was added slowly to the resulting clarified cell extract with stirring at 4°C to achieve 50% saturation. After 1 hour, the mixture was centrifuged at 23,000 x g for 30 minutes. The supernatant was then brought to 85% saturation by the addition of solid ammonium sulfate (159 g/liter) and stirred for 1h at 4°C before centrifugation (23,000 xg for 30 min). The resultant 50-85% ammonium sulfate pellet was resuspended in 600 mL of Lysing Buffer and the residual ammonium sulfate was removed by dialysis against the same buffer at 4°C. The desalted material was centrifuged twice to remove particulate matter (23,000 xg for 30 min) and 700 - 800 Units of the clarified material was loaded onto a Red-120 dye affinity column (32 mm X 140 mm) equilibrated in 50 mM Tris-Cl pH 7.5, 1 mM MgCl₂, pepstatin (1 µg/mL), leupeptin (1.25 µg/mL), and aprotinin (2.5 µg/mL). Reductase activity was eluted from the column at a flowrate of 8 mL/min under the following conditions:

- 1) a 10 minute linear gradient from 0 - 0.3 M NaCl; 2) 13 minutes at 0.3 M NaCl; 3) a 60 minute linear gradient from 0.3 - 1.5 M NaCl. The fractions containing reductase activity were pooled, and changed to 20 mM potassium phosphate buffer (pH 7.2), pepstatin (1 µg/mL), leupeptin (1.25 µg/mL), and aprotinin (2.5 µg/mL) by dialysis at 4°C.

The sample was clarified by centrifugation (23,000 x g for 30 min) and 400 Units was loaded onto a Bio-Scale CHT-I hydroxyapatite column (15 mm x 113 mm, Bio-Rad, Inc.) equilibrated in the same buffer that had been made 5% in glycerol. Reductase activity was eluted from the column at

-25-

a flowrate of 5.0 mL/min in a sodium chloride step gradient consisting of 5 minutes at 0 M NaCl, a gradient step to 0.7 M NaCl which was maintained for 10 minutes, and then a 20 minute linear gradient from 0.7 - 1.0 M NaCl. The fractions
5 containing reductase activity were pooled and desalted with 20 mM potassium phosphate buffer (pH 7.2), pepstatin A (1 μ g/mL), leupeptin (1.25 μ g/mL), and aprotinin (2.5 μ g/mL) by dialysis at 4°C. The sample (100- 200 Units) was loaded onto a Bio-Scale CHT-I hydroxyapatite column (10 mm x 64 mm)
10 equilibrated in the same buffer which had been made 5% in glycerol. Reductase activity was eluted from the column at a flowrate of 2.0 mL/min in a 25 minute linear gradient from 0 to 50% 400 mM potassium phosphate (pH 6.8), 5% glycerol. Fractions containing reductase activity were pooled and
15 changed into 10 mM Tris-Cl (pH 8.5) by dialysis at 4°C. The sample was then made 10% in glycerol, concentrated to 0.4 mg/mL by ultrafiltration (Amicon, YM-10), and stored at -70°C.

20

EXAMPLE 3

Reductase Activity Using the Ketoreductase from *Z. rouxii*

Reductase activity was measured using a suitable substrate and a partially purified or substantially purified ketoreductase from *Z. rouxii*. Activity was measured as a
25 function of the absorbance change at 340 nm, resulting from the oxidation of NADPH. The 1 ml assay contained a mixture of 3.0 mM 3,4-methylenedioxyphenyl acetone, 162 μ M NADPH, 50 mM MOPS buffer (pH 6.8), and 0.6 mU of ketoreductase and was carried out at 26° C. Reaction mixtures were first
30 equilibrated at 26°C for 10 min in the absence of NADPH, and

-26-

then initiated by addition of NADPH. The absorbance was measured at 340 nm every 15 seconds over a 5 minute period; the change in absorbance was found to be linear over that time period. The kinetic parameters for 3,4-methylenedioxyphenyl acetone were determined at an NADPH concentration of 112 μM and a 3,4-methylenedioxyphenyl acetone concentration that varied from 1.7 mM - 7.2 mM. The kinetic parameters for NADPH were determined by maintaining the 3,4-methylenedioxyphenyl acetone concentration at 3 mM and the NADPH concentration was varied from 20.5 μM - 236.0 μM . An extinction coefficient of 6220 $\text{M}^{-1} \text{cm}^{-1}$ for NADPH absorbance at 340 nm was used to calculate the specific activity of the enzyme. For assays using isatin, the change in absorbance with time was measured at 414 nm using an extinction coefficient of 849 $\text{M}^{-1} \text{cm}^{-1}$ to calculate activity. One Unit of activity corresponds to 1 μmol of NADPH consumed per minute. For assays carried out at differing pH values, 10 mM Bis-Tris and 10 mM Tris were adjusted to the appropriate pH with HCl. Kinetic parameters were determined by non-linear regression using the JMP[®] statistics and graphics program.

EXAMPLE 4

Whole Cell Method for Stereoselective Reduction of Ketone

Using Recombinant Yeast Cell

A vector for expressing the cloned *Z. rouxii* ketoreductase gene (SEQ ID NO:1) in a procaryotic or fungal cell, such as *S. cerevisiae*, is constructed as follows. A 1014 base pair fragment of *Z. rouxii* genomic DNA or cDNA, carrying the ketoreductase gene, is amplified by PCR using

-27-

primers targeted to the ends of the coding region specified in SEQ ID NO:1. It is desirable that the primers also incorporate suitable cloning sites for cloning of said 1014 base pair fragment into an expression vector. The

5 appropriate fragment encoding ketoreductase is amplified and purified using standard methods, for cloning into an expression vector.

A suitable vector for expression in *E. coli* and *S. cerevisiae* is pYX213 (available from Novagen, Inc., 597
10 Science Drive, Madison, WI 53711; Code MBV-029-10), a 7.5 Kb plasmid that carries the following genetic markers: ori, 2 μ circle, Amp^R, CEN, URA3, and the GAL promoter, for high level expression in yeast. Downstream of the GAL promoter, pYX213 carries a multiple cloning site (MCS), which will
15 accommodate the ketoreductase gene amplified in the preceding step. A recombinant plasmid is created by digesting pYX213 and the amplified ketoreductase gene with a restriction enzyme, such as BamH1, and ligating the fragments together.

20 A recombinant expression vector carrying the *Z. rouxii* ketoreductase gene is transformed into a suitable Ura⁻ strain of *S. cerevisiae*, using well known methods. Ura⁺ transformants are selected on minimal medium lacking uracil.

Expression of the recombinant ketoreductase gene
25 may be induced if desired by growing transformants in minimal medium that contains 2% galactose as the sole carbon source.

To carry out a whole cell stereospecific reduction, 3,4-methylenedioxyphenyl acetone is added to a
30 culture of transformants to a concentration of about 10

-28-

grams per liter of culture. The culture is incubated with shaking at room temperature for 24 hours, and the presence of the chiral alcohol analyzed by HPLC.

-29-

WE CLAIM:

1. A substantially pure ketoreductase protein having the amino acid sequence which is SEQ ID NO:2.
- 5 2. An isolated nucleic acid compound encoding the protein of Claim 1, said protein having the amino acid sequence which is SEQ ID NO:2.
- 10 3. An isolated nucleic acid compound encoding the protein of Claim 1, wherein said compound has a sequence selected from the group consisting of:
 - (a) SEQ ID NO:1; or
 - (b) SEQ ID NO:3.
- 15 4. An isolated nucleic acid compound of Claim 3 wherein the sequence of said compound is SEQ ID NO:1
5. An isolated nucleic acid compound having a sequence
- 20 complementary to SEQ ID NO:1.
6. An isolated nucleic acid compound of Claim 3 wherein the sequence of said compound is SEQ ID NO:3.
- 25 7. An isolated nucleic acid compound having a sequence complementary to SEQ ID NO:3.
8. A vector comprising an isolated nucleic acid compound of Claim 2.
- 30 9. A vector comprising an isolated nucleic acid compound of Claim 3.
10. A vector of Claim 9, wherein said isolated nucleic acid
- 35 compound is SEQ ID NO:1 operably-linked to a promoter sequence.

-30-

11. A host cell containing the vector of Claim 10.

12. A method for constructing a recombinant host cell having
5 the potential to express SEQ ID NO:2, said method comprising
introducing into said host cell by any suitable means a
vector of Claim 9.

13. A method for expressing SEQ ID NO:2 in the recombinant
10 host cell of Claim 12, said method comprising culturing said
recombinant host cell under conditions suitable for gene
expression.

14. A method for reducing a ketone in a stereospecific
15 manner comprising providing a quantity of a suitable ketone
to a culture of recombinant cells for a suitable period of
time, wherein said cells are transformed with a vector that
carries a ketoreductase gene, and wherein said cells express
said ketoreductase gene.

20

15. A method, as in claim 14 wherein said gene is selected
from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ
ID NO:7, SEQ ID NO:10, and SEQ ID NO:13.

25 16. A method, as in claim 14 wherein said ketone comprises
an α -ketolactone, α -ketolactam, or a diketone.

17. A method, as in Claim 14, wherein said recombinant cells
are selected from the group consisting of *S. cerevisiae*, *Z.*
30 *rouxii*, and *E. coli*.

18. A method for reducing a ketone in a stereospecific
manner comprising mixing a quantity of a suitable ketone
with a substantially purified ketoreductase and suitable
35 reducing agent.

-31-

19. A method, as in Claim 18 wherein said ketoreductase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:8, SEQ ID NO:11, and SEQ ID NO:14.

5 20. An isolated nucleic acid compound that encodes a protein having ketoreductase activity wherein said nucleic acid hybridizes under high stringency conditions to SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:7, SEQ ID NO:10, or SEQ ID NO:13.

10 21. A method, as in Claim 18 wherein said reducing agent is NADPH.

- 1 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Costello, Colleen A.
Menke, Michael A.
Hershberger, Charles L.
Zmijewski, Milton J.
- (ii) TITLE OF INVENTION: Ketoreductase Gene and Protein From Yeast
- (iii) NUMBER OF SEQUENCES: 15
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Eli Lilly and Company
 - (B) STREET: Lilly Corporate Center
 - (C) CITY: Indianapolis
 - (D) STATE: Indiana
 - (E) COUNTRY: United States
 - (F) ZIP: 46285
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Webster, Thomas D.
 - (B) REGISTRATION NUMBER: 39,872
 - (C) REFERENCE/DOCKET NUMBER: X-11325
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 317-276-3334

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1270 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO

- 2 -

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 164..1177

(D) OTHER INFORMATION: Z.rouxii ketoreductase

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGAATGGTTA TTTTAGCAAT TGCTGTGTGA GGCAC TGACC TAAAGATGTG TATAAATAGT	60
GGGACTGTGT ACTCATGAGG ATCAATACAT GTATAAACTT ACCATACTTT CACACAAGTC	120
AACTTAGAAT CAATCAATCA ATCAATTAAT CAAGCTATAC AAT ATG ACA AAA GTC	175
Met Thr Lys Val	
1	
TTC GTA ACA GGT GCC AAC GGA TTC GTT GCT CAA CAC GTC GTT CAT CAA	223
Phe Val Thr Gly Ala Asn Gly Phe Val Ala Gln His Val Val His Gln	
5 10 15 20	
CTA TTA GAA AAG AAC TAT ACA GTG GTT GGA TCT GTC CGT TCA ACT GAG	271
Leu Leu Glu Lys Asn Tyr Thr Val Val Gly Ser Val Arg Ser Thr Glu	
25 30 35	
AAA GGT GAT AAA TTA GCT AAA TTG CTA AAC AAT CCA AAA TTT TCA TAT	319
Lys Gly Asp Lys Leu Ala Lys Leu Leu Asn Asn Pro Lys Phe Ser Tyr	
40 45 50	
GAG ATT ATT AAA GAT ATG GTC AAT TCG AGA GAT GAA TTC GAT AAG GCT	367
Glu Ile Ile Lys Asp Met Val Asn Ser Arg Asp Glu Phe Asp Lys Ala	
55 60 65	
TTA CAA AAA CAT TCA GAT GTT GAA ATT GTC TTA CAT ACT GCT TCA CCA	415
Leu Gln Lys His Ser Asp Val Glu Ile Val Leu His Thr Ala Ser Pro	
70 75 80	
GTC TTC CCA GGT GGT ATT AAA GAT GTT GAA AAA GAA ATG ATC CAA CCA	463
Val Phe Pro Gly Gly Ile Lys Asp Val Glu Lys Glu Met Ile Gln Pro	
85 90 95 100	
GCT GTT AAT GGT ACT AGA AAT GTC TTG TTA TCA ATC AAG GAT AAC TTA	511
Ala Val Asn Gly Thr Arg Asn Val Leu Leu Ser Ile Lys Asp Asn Leu	
105 110 115	
CCA AAT GTC AAG AGA TTT GTT TAC ACT TCT TCA TTA GCT GCT GTC CGT	559
Pro Asn Val Lys Arg Phe Val Tyr Thr Ser Ser Leu Ala Ala Val Arg	
120 125 130	
ACT GAA GGT GCT GGT TAT AGT GCA GAC GAA GTT GTC ACC GAA GAT TCT	607
Thr Glu Gly Ala Gly Tyr Ser Ala Asp Glu Val Val Thr Glu Asp Ser	
135 140 145	
TGG AAC AAT ATT GCA TTG AAA GAT GCC ACC AAG GAT GAA GGT ACA GCT	655
Trp Asn Asn Ile Ala Leu Lys Asp Ala Thr Lys Asp Glu Gly Thr Ala	

- 3 -

150	155	160	
TAT GAG GCT TCC AAG ACA	TAT GGT GAA AAA GAA	GTT TGG AAT TTC TTC	703
Tyr Glu Ala Ser Lys Thr	Tyr Gly Glu Lys Glu	Val Trp Asn Phe Phe	
165	170	175 180	
GAA AAA ACT AAA AAT GTT AAT TTC	GAT TTT GCC ATC ATC AAC CCA GTT		751
Glu Lys Thr Lys Asn Val Asn Phe Asp	Phe Ala Ile Ile Asn Pro Val		
185	190	195	
TAT GTC TTT GGT CCT CAA TTA TTT	GAA GAA TAC GTT ACT GAT AAA TTG		799
Tyr Val Phe Gly Pro Gln Leu Phe	Glu Glu Tyr Val Thr Asp Lys Leu		
200	205	210	
AAC TTT TCC AGT GAA ATC ATT AAT AGT	ATA ATA AAA GGT GAA AAG AAG		847
Asn Phe Ser Ser Glu Ile Ile Asn Ser	Ile Ile Lys Gly Glu Lys Lys		
215	220	225	
GAA ATT GAA GGT TAT GAA ATT GAT GTT	AGA GAT ATT GCA AGA GCT CAT		895
Glu Ile Glu Gly Tyr Glu Ile Asp Val	Arg Asp Ile Ala Arg Ala His		
230	235	240	
ATC TCT GCT GTT GAA AAT CCA GCA ACT	ACA CGT CAA AGA TTA ATT CCA		943
Ile Ser Ala Val Glu Asn Pro Ala Thr	Thr Arg Gln Arg Leu Ile Pro		
245	250	255 260	
GCA GTT GCA CCA TAC AAT CAA CAA ACT	ATC TTG GAT GTT TTG AAT GAA		991
Ala Val Ala Pro Tyr Asn Gln Gln Thr	Ile Leu Asp Val Leu Asn Glu		
265	270	275	
AAC TTC CCA GAA TTG AAA GGT AAA ATC	GAT GTT GGG AAA CCA GGT TCT		1039
Asn Phe Pro Glu Leu Lys Gly Lys Ile	Asp Val Gly Lys Pro Gly Ser		
280	285	290	
CAA AAT GAA TTT ATT AAA AAA TAT TAT	AAA TTA GAT AAC TCA AAG ACC		1087
Gln Asn Glu Phe Ile Lys Lys Tyr Tyr	Lys Leu Asp Asn Ser Lys Thr		
295	300	305	
AAA AAA GTT TTA GGT TTT GAA TTC ATT	TCC CAA GAG CAA ACA ATC AAA		1135
Lys Lys Val Leu Gly Phe Glu Phe Ile	Ser Gln Glu Gln Thr Ile Lys		
310	315	320	
GAT GCT GCT GCT CAA ATC TTG TCC GTT	AAA AAT GGA AAA AAA		1177
Asp Ala Ala Ala Gln Ile Leu Ser Val	Lys Asn Gly Lys Lys		
325	330	335	
TAAGTGAACT AGACCTGTCA CTATCAGATT	ATTAGAGTTC TGTATAGATT AAAGTGTGAA		1237
AATGTATTAG AATCATAATT TTATAATATG	CCT		1270

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 338 amino acids

- 4 -

(B) TYPE: amino acid
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

Met Thr Lys Val Phe Val Thr Gly Ala Asn Gly Phe Val Ala Gln His
 1             5             10             15

Val Val His Gln Leu Leu Glu Lys Asn Tyr Thr Val Val Gly Ser Val
      20             25             30

Arg Ser Thr Glu Lys Gly Asp Lys Leu Ala Lys Leu Leu Asn Asn Pro
      35             40             45

Lys Phe Ser Tyr Glu Ile Ile Lys Asp Met Val Asn Ser Arg Asp Glu
 50             55             60

Phe Asp Lys Ala Leu Gln Lys His Ser Asp Val Glu Ile Val Leu His
 65             70             75             80

Thr Ala Ser Pro Val Phe Pro Gly Gly Ile Lys Asp Val Glu Lys Glu
      85             90             95

Met Ile Gln Pro Ala Val Asn Gly Thr Arg Asn Val Leu Leu Ser Ile
      100             105             110

Lys Asp Asn Leu Pro Asn Val Lys Arg Phe Val Tyr Thr Ser Ser Leu
      115             120             125

Ala Ala Val Arg Thr Glu Gly Ala Gly Tyr Ser Ala Asp Glu Val Val
      130             135             140

Thr Glu Asp Ser Trp Asn Asn Ile Ala Leu Lys Asp Ala Thr Lys Asp
      145             150             155             160

Glu Gly Thr Ala Tyr Glu Ala Ser Lys Thr Tyr Gly Glu Lys Glu Val
      165             170             175

Trp Asn Phe Phe Glu Lys Thr Lys Asn Val Asn Phe Asp Phe Ala Ile
      180             185             190

Ile Asn Pro Val Tyr Val Phe Gly Pro Gln Leu Phe Glu Glu Tyr Val
      195             200             205

Thr Asp Lys Leu Asn Phe Ser Ser Glu Ile Ile Asn Ser Ile Ile Lys
      210             215             220

Gly Glu Lys Lys Glu Ile Glu Gly Tyr Glu Ile Asp Val Arg Asp Ile
      225             230             235             240

Ala Arg Ala His Ile Ser Ala Val Glu Asn Pro Ala Thr Thr Arg Gln
      245             250             255

```

- 5 -

Arg Leu Ile Pro Ala Val Ala Pro Tyr Asn Gln Gln Thr Ile Leu Asp
 260 265 270

Val Leu Asn Glu Asn Phe Pro Glu Leu Lys Gly Lys Ile Asp Val Gly
 275 280 285

Lys Pro Gly Ser Gln Asn Glu Phe Ile Lys Lys Tyr Tyr Lys Leu Asp
 290 295 300

Asn Ser Lys Thr Lys Lys Val Leu Gly Phe Glu Phe Ile Ser Gln Glu
 305 310 315 320

Gln Thr Ile Lys Asp Ala Ala Ala Gln Ile Leu Ser Val Lys Asn Gly
 325 330 335

Lys Lys

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1271 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

UGAAUGGUUA UUUUAGCAAU UGCUGUGUGA GGCACUGACC UAAAGAUGUG UAUAAAUAGU	60
GGGACUGUGU ACUCAUGAGG AUCAAUACAU GUAAUAAACUU ACCAUACUUU CACACAAGUC	120
AACUUAGAAU CAAUCAAUCA AUCAAUUAAU CAAGCUAUAC AAUAUGACAA AAGUCUUCGU	180
AACAGGUGCC AACGGAUUCG UUGCUCACAA CGUCGUUCAU CAACUAUUAG AAAAGAACUA	240
UACAGUGGUU GGAUCUGUCC GUUCAACUGA GAAAGGUGAU AAUUUAGCUA AAUUGCUAAA	300
CAAUCCAAAA UUUUCAUAUG AGAUUAUUAA AGAUUAGGUC AAUUCGAGAG AUGAAUUCGA	360
UAAGGCUUUA CAAAAACAUU CAGAUGUUGA AAUUGUCUUA CAUACUGCUU CACCAGUCUU	420
CCCAGGUGGU AUUAAAGAUG UUGAAAAAGA AAUGAUCCAA CCAGCUGUUA AUGGUACUAG	480
AAAUGUCUUG UUAUCAAUCA AGGAUAACUU ACCAAAUGUC AAGAGAUUUG UUUACACUUC	540

- 6 -

```

UUCAUUAGCTU GCUGUCCGUA CUGAAGGUGC UGGUUUAUAGU GCAGACGAAG UUGUCACCGA      600
AGAUUCUUGG AACAAUAUUG CAUUGAAAGA UGCCACCAAG GAUGAAGGUA CAGCUUAUGA      660
GGCUUCCAAG ACAUAUGGUG AAAAAGAAGU UUGGAAUUTUC UUCGAAAAAA CUAAAAAUGU      720
UAAUUUCGAU UUUGCCAUCA UCAACCCAGU UUAUGUCUUU GGUCCUCAAU UAUUUGAAGA      780
AUACGUUACU GAUAAAUUGA ACUUUUCCAG UGAAAUCAUU AAUAGUAUAA UAAAAGGUGA      840
AAAGAAGGAA AUUGAAGGUU AUGAAAUUGA UGUUAGAGAU AUUGCAAGAG CUCAUAUCUC      900
UGCUGUUGAA AAUCCAGCAA CUACACGUCA AAGAUUAAUU CCAGCAGUUG CACCAUACAA      960
UCAACAAACU AUCUUGGAUG UUUUGAAUGA AAACUUCCCA GAAUUGAAAAG GUAAAAUCGA     1020
UGUUGGGAAA CCAGGUUCUC AAAAUGAAUU UAUAUAAAAA UAUAUAAAAU UAGAUAAACUC     1080
AAAGACCAAA AAAGUUUUAG GUUUUGAAUU CAUUUCCCAA GAGCAAACAA UCAAAGAUGC     1140
UGCUGCUCAA AUCUUGUCCG UUA AAAAUGG AAAAAAUAA GUGAACUAGA CCUGUCACUA     1200
UCAGAUUAUU AGAGUUCUGU AUAGAUUAAA GUGUGAAAAU GUAAUAGAAU CAUAAUUUUA     1260
UAAUUAUGCC U                                                                1271

```

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1032 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1032
- (D) OTHER INFORMATION: *S.cerevisiae* YDR541c

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

ATG TCT AAT ACA GTT CTA GTT TCT GGC GCT TCA GGT TTT ATT GCC TTG      48
Met Ser Asn Thr Val Leu Val Ser Gly Ala Ser Gly Phe Ile Ala Leu
  1             5             10             15

CAT ATC CTG TCA CAA TTG TTA AAA CAA GAT TAT AAG GTT ATT GGA ACT      96
His Ile Leu Ser Gln Leu Leu Lys Gln Asp Tyr Lys Val Ile Gly Thr
      20             25             30

```

- 7 -

GTG AGA TCC CAT GAA AAA GAA GCA AAA TTG CTA AGA CAA TTT CAA CAT	144
Val Arg Ser His Glu Lys Glu Ala Lys Leu Leu Arg Gln Phe Gln His	
35 40 45	
AAC CCT AAT TTA ACT TTA GAA ATT GTT CCG GAC ATT TCT CAT CCA AAT	192
Asn Pro Asn Leu Thr Leu Glu Ile Val Pro Asp Ile Ser His Pro Asn	
50 55 60	
GCT TTC GAT AAG GTT CTG CAG AAA CGT GGA CGT GAG ATT AGG TAT GTT	240
Ala Phe Asp Lys Val Leu Gln Lys Arg Gly Arg Glu Ile Arg Tyr Val	
65 70 75 80	
CTA CAC ACG GCC TCT CCT TTT CAT TAT GAT ACT ACC GAA TAT GAA AAA	288
Leu His Thr Ala Ser Pro Phe His Tyr Asp Thr Thr Glu Tyr Glu Lys	
85 90 95	
GAC TTA TTG ATT CCC GCG TTA GAA GGT ACA AAA AAC ATC CTA AAT TCT	336
Asp Leu Leu Ile Pro Ala Leu Glu Gly Thr Lys Asn Ile Leu Asn Ser	
100 105 110	
ATC AAG AAA TAT GCA GCA GAC ACT GTA GAG CGT GTT GTT GTG ACT TCT	384
Ile Lys Lys Tyr Ala Ala Asp Thr Val Glu Arg Val Val Val Thr Ser	
115 120 125	
TCT TGT ACT GCT ATT ATA ACC CTT GCA AAG ATG GAC GAT CCC AGT GTG	432
Ser Cys Thr Ala Ile Ile Thr Leu Ala Lys Met Asp Asp Pro Ser Val	
130 135 140	
GTT TTT ACA GAA GAG AGT TGG AAC GAA GCA ACC TGG GAA AGC TGT CAA	480
Val Phe Thr Glu Glu Ser Trp Asn Glu Ala Thr Trp Glu Ser Cys Gln	
145 150 155 160	
ATT GAT GGG ATA AAT GCT TAC TTT GCA TCC AAG AAG TTT GCT GAA AAG	528
Ile Asp Gly Ile Asn Ala Tyr Phe Ala Ser Lys Lys Phe Ala Glu Lys	
165 170 175	
GCT GCC TGG GAG TTC ACA AAA GAG AAT GAA GAT CAC ATC AAA TTC AAA	576
Ala Ala Trp Glu Phe Thr Lys Glu Asn Glu Asp His Ile Lys Phe Lys	
180 185 190	
CTA ACA ACA GTC AAC CCT TCT CTT CTT TTT GGT CCT CAA CTT TTC GAT	624
Leu Thr Thr Val Asn Pro Ser Leu Leu Phe Gly Pro Gln Leu Phe Asp	
195 200 205	
GAA GAT GTG CAT GGC CAT TTG AAT ACT TCT TGC GAA ATG ATC AAT GGC	672
Glu Asp Val His Gly His Leu Asn Thr Ser Cys Glu Met Ile Asn Gly	
210 215 220	
CTA ATT CAT ACC CCA GTA AAT GCC AGT GTT CCT GAT TTT CAT TCC ATT	720
Leu Ile His Thr Pro Val Asn Ala Ser Val Pro Asp Phe His Ser Ile	
225 230 235 240	
TTT ATT GAT GTA AGG GAT GTG GCC CTA GCT CAT CTG TAT GCT TTC CAG	768
Phe Ile Asp Val Arg Asp Val Ala Leu Ala His Leu Tyr Ala Phe Gln	

- 8 -

245	250	255	
AAG GAA AAT ACC GCG GGT AAA AGA TTA GTG GTA ACT AAC GGT AAA TTT			816
Lys Glu Asn Thr Ala Gly Lys Arg Leu Val Val Thr Asn Gly Lys Phe			
260	265	270	
GGA AAC CAA GAT ATC CTG GAT ATT TTG AAC GAA GAT TTT CCA CAA TTA			864
Gly Asn Gln Asp Ile Leu Asp Ile Leu Asn Glu Asp Phe Pro Gln Leu			
275	280	285	
AGA GGT CTC ATT CCT TTG GGT AAG CCT GGC ACA GGT GAT CAA GTC ATT			912
Arg Gly Leu Ile Pro Leu Gly Lys Pro Gly Thr Gly Asp Gln Val Ile			
290	295	300	
GAC CGC GGT TCA ACT ACA GAT AAT AGT GCA ACG AGG AAA ATA CTT GGC			960
Asp Arg Gly Ser Thr Thr Asp Asn Ser Ala Thr Arg Lys Ile Leu Gly			
305	310	315	320
TTT GAG TTC AGA AGT TTA CAC GAA AGT GTC CAT GAT ACT GCT GCC CAA			1008
Phe Glu Phe Arg Ser Leu His Glu Ser Val His Asp Thr Ala Ala Gln			
325	330	335	
ATT TTG AAG AAG GAG AAC AGA TTA			1032
Ile Leu Lys Lys Glu Asn Arg Leu			
340			

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 344 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met	Ser	Asn	Thr	Val	Leu	Val	Ser	Gly	Ala	Ser	Gly	Phe	Ile	Ala	Leu	
1				5				10					15			
His	Ile	Leu	Ser	Gln	Leu	Leu	Lys	Gln	Asp	Tyr	Lys	Val	Ile	Gly	Thr	
		20					25						30			
Val	Arg	Ser	His	Glu	Lys	Glu	Ala	Lys	Leu	Leu	Arg	Gln	Phe	Gln	His	
		35					40					45				
Asn	Pro	Asn	Leu	Thr	Leu	Glu	Ile	Val	Pro	Asp	Ile	Ser	His	Pro	Asn	
		50				55				60						
Ala	Phe	Asp	Lys	Val	Leu	Gln	Lys	Arg	Gly	Arg	Glu	Ile	Arg	Tyr	Val	
		65				70				75				80		
Leu	His	Thr	Ala	Ser	Pro	Phe	His	Tyr	Asp	Thr	Thr	Glu	Tyr	Glu	Lys	
					85				90					95		

- 9 -

Asp Leu Leu Ile Pro Ala Leu Glu Gly Thr Lys Asn Ile Leu Asn Ser
 100 105 110
 Ile Lys Lys Tyr Ala Ala Asp Thr Val Glu Arg Val Val Val Thr Ser
 115 120 125
 Ser Cys Thr Ala Ile Ile Thr Leu Ala Lys Met Asp Asp Pro Ser Val
 130 135 140
 Val Phe Thr Glu Glu Ser Trp Asn Glu Ala Thr Trp Glu Ser Cys Gln
 145 150 155 160
 Ile Asp Gly Ile Asn Ala Tyr Phe Ala Ser Lys Lys Phe Ala Glu Lys
 165 170 175
 Ala Ala Trp Glu Phe Thr Lys Glu Asn Glu Asp His Ile Lys Phe Lys
 180 185 190
 Leu Thr Thr Val Asn Pro Ser Leu Leu Phe Gly Pro Gln Leu Phe Asp
 195 200 205
 Glu Asp Val His Gly His Leu Asn Thr Ser Cys Glu Met Ile Asn Gly
 210 215 220
 Leu Ile His Thr Pro Val Asn Ala Ser Val Pro Asp Phe His Ser Ile
 225 230 235 240
 Phe Ile Asp Val Arg Asp Val Ala Leu Ala His Leu Tyr Ala Phe Gln
 245 250 255
 Lys Glu Asn Thr Ala Gly Lys Arg Leu Val Val Thr Asn Gly Lys Phe
 260 265 270
 Gly Asn Gln Asp Ile Leu Asp Ile Leu Asn Glu Asp Phe Pro Gln Leu
 275 280 285
 Arg Gly Leu Ile Pro Leu Gly Lys Pro Gly Thr Gly Asp Gln Val Ile
 290 295 300
 Asp Arg Gly Ser Thr Thr Asp Asn Ser Ala Thr Arg Lys Ile Leu Gly
 305 310 315 320
 Phe Glu Phe Arg Ser Leu His Glu Ser Val His Asp Thr Ala Ala Gln
 325 330 335
 Ile Leu Lys Lys Glu Asn Arg Leu
 340

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1032 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

- 10 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

AUGUCUAAUA CAGUUCUAGU UUCUGGCGCU UCAGGUUUUA UUGCCUUGCA UAUCCUGUCA	60
CAAUGUUUAA AACAAAGAUUA UAAGGUUAUU GGAACUGUGA GAUCCCAUGA AAAAGAAGCA	120
AAAUUGCUAA GACAAUUUCA ACAUAACCCU AAUUUAACTU UAGAAAUUGU UCCGGACAUU	180
UCUCAUCCAA AUGCUUUCGA UAAGGUUCUG CAGAAACGUG GACGUGAGAU UAGGUAUGUU	240
CUACACACGG CCUCUCCUUU UCAUUAUGAU ACUACCGAAU AUGAAAAAGA CUUUAUGAUU	300
CCCGCGUUAG AAGGUACAAA AAACAUCCUA AAUUCUAUCA AGAAAUAGC AGCAGACACU	360
GUAGAGCGUG UUGUUGUGAC UUCUUCUUGU ACUGCUAUUA UAACCCUUGC AAAGAUGGAC	420
GAUCCCAGUG UGGUUUUUAC AGAAGAGAGU UGGAACGAAG CAACCUUGGA AAGCUGUCAA	480
AUUGAUGGGA UAAAUUGCUUA CUUUGCAUCC AAGAAGUUUG CUGAAAAGGC UGCCUGGGAG	540
UUCACAAAAG AGAAUGAAGA UCACAUCAA UUCAACUAA CAACAGUCA CCCUUCUCUU	600
CUUUUUGGUC CUCAACUUUU CGAUGAAGAU GUGCAUGGCC AUUUGAAUAC UUCUUGCGAA	660
AUGAUCAAUG GCCUAAUUA UACCCCAGUA AAUGCCAGUG UCCUGAUUU UCAUUCCAUU	720
UUUAUUGAUG UAAGGGAUGU GGCCCUAGCU CAUCUGUAUG CUUCCAGAA GGAAAUUACC	780
GCGGGUAAAA GAUUGUGGU AACUAACGGU AAUUUUGGAA ACCAAGAUU CCUGGAUUAU	840
UGAACGAAG AUUUUCCACA AUUAAGAGGU CUCAUUCUU UGGGUAAGCC UGGCACAGGU	900
GAUCAAGUCA UUGACCGCGG UUCAACUACA GAUAAUAGUG CAACGAGGAA AAUACUUGGC	960
UUUGAGUUA GAAGUUUACA CGAAAGUGUC CAUGAUACUG CUGCCCAAU UUUGAAGAAG	1020
GAGAACAGAU UA	1032

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1029 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

- 11 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1026

(D) OTHER INFORMATION: S.cerevisiae YOL151W

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATG TCA GTT TTC GTT TCA GGT GCT AAC GGG TTC ATT GCC CAA CAC ATT	48
Met Ser Val Phe Val Ser Gly Ala Asn Gly Phe Ile Ala Gln His Ile	
1 5 10 15	
GTC GAT CTC CTG TTG AAG GAA GAC TAT AAG GTC ATC GGT TCT GCC AGA	96
Val Asp Leu Leu Leu Lys Glu Asp Tyr Lys Val Ile Gly Ser Ala Arg	
20 25 30	
AGT CAA GAA AAG GCC GAG AAT TTA ACG GAG GCC TTT GGT AAC AAC CCA	144
Ser Gln Glu Lys Ala Glu Asn Leu Thr Glu Ala Phe Gly Asn Asn Pro	
35 40 45	
AAA TTC TCC ATG GAA GTT GTC CCA GAC ATA TCT AAG CTG GAC GCA TTT	192
Lys Phe Ser Met Glu Val Val Pro Asp Ile Ser Lys Leu Asp Ala Phe	
50 55 60	
GAC CAT GTT TTC CAA AAG CAC GGC AAG GAT ATC AAG ATA GTT CTA CAT	240
Asp His Val Phe Gln Lys His Gly Lys Asp Ile Lys Ile Val Leu His	
65 70 75 80	
ACG GCC TCT CCA TTC TGC TTT GAT ATC ACT GAC AGT GAA CGC GAT TTA	288
Thr Ala Ser Pro Phe Cys Phe Asp Ile Thr Asp Ser Glu Arg Asp Leu	
85 90 95	
TTA ATT CCT GCT GTG AAC GGT GTT AAG GGA ATT CTC CAC TCA ATT AAA	336
Leu Ile Pro Ala Val Asn Gly Val Lys Gly Ile Leu His Ser Ile Lys	
100 105 110	
AAA TAC GCC GCT GAT TCT GTA GAA CGT GTA GTT CTC ACC TCT TCT TAT	384
Lys Tyr Ala Ala Asp Ser Val Glu Arg Val Val Leu Thr Ser Ser Tyr	
115 120 125	
GCA GCT GTG TTC GAT ATG GCA AAA GAA AAC GAT AAG TCT TTA ACA TTT	432
Ala Ala Val Phe Asp Met Ala Lys Glu Asn Asp Lys Ser Leu Thr Phe	
130 135 140	
AAC GAA GAA TCC TGG AAC CCA GCT ACC TGG GAG AGT TGC CAA AGT GAC	480
Asn Glu Glu Ser Trp Asn Pro Ala Thr Trp Glu Ser Cys Gln Ser Asp	
145 150 155 160	

- 12 -

CCA GTT AAC GCC TAC TGT GGT TCT AAG AAG TTT GCT GAA AAA GCA GCT	528
Pro Val Asn Ala Tyr Cys Gly Ser Lys Lys Phe Ala Glu Lys Ala Ala	
165 170 175	
TGG GAA TTT CTA GAG GAG AAT AGA GAC TCT GTA AAA TTC GAA TTA ACT	576
Trp Glu Phe Leu Glu Glu Asn Arg Asp Ser Val Lys Phe Glu Leu Thr	
180 185 190	
GCC GTT AAC CCA GTT TAC GTT TTT GGT CCG CAA ATG TTT GAC AAA GAT	624
Ala Val Asn Pro Val Tyr Val Phe Gly Pro Gln Met Phe Asp Lys Asp	
195 200 205	
GTG AAA AAA CAC TTG AAC ACA TCT TGC GAA CTC GTC AAC AGC TTG ATG	672
Val Lys Lys His Leu Asn Thr Ser Cys Glu Leu Val Asn Ser Leu Met	
210 215 220	
CAT TTA TCA CCA GAG GAC AAG ATA CCG GAA CTA TTT GGT GGA TAC ATT	720
His Leu Ser Pro Glu Asp Lys Ile Pro Glu Leu Phe Gly Gly Tyr Ile	
225 230 235 240	
GAT GTT CGT GAT GTT GCA AAG GCT CAT TTA GTT GCC TTC CAA AAG AGG	768
Asp Val Arg Asp Val Ala Lys Ala His Leu Val Ala Phe Gln Lys Arg	
245 250 255	
GAA ACA ATT GGT CAA AGA CTA ATC GTA TCG GAG GCC AGA TTT ACT ATG	816
Glu Thr Ile Gly Gln Arg Leu Ile Val Ser Glu Ala Arg Phe Thr Met	
260 265 270	
CAG GAT GTT CTC GAT ATC CTT AAC GAA GAC TTC CCT GTT CTA AAA GGC	864
Gln Asp Val Leu Asp Ile Leu Asn Glu Asp Phe Pro Val Leu Lys Gly	
275 280 285	
AAT ATT CCA GTG GGG AAA CCA GGT TCT GGT GCT ACC CAT AAC ACC CTT	912
Asn Ile Pro Val Gly Lys Pro Gly Ser Gly Ala Thr His Asn Thr Leu	
290 295 300	
GGT GCT ACT CTT GAT AAT AAA AAG AGT AAG AAA TTG TTA GGT TTC AAG	960
Gly Ala Thr Leu Asp Asn Lys Lys Ser Lys Lys Leu Leu Gly Phe Lys	
305 310 315 320	
TTC AGG AAC TTG AAA GAG ACC ATT GAC GAC ACT GCC TCC CAA ATT TTA	1008
Phe Arg Asn Leu Lys Glu Thr Ile Asp Asp Thr Ala Ser Gln Ile Leu	
325 330 335	
AAA TTT GAG GGC AGA ATA TAA	1029
Lys Phe Glu Gly Arg Ile	
340	

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 342 amino acids

(B) TYPE: amino acid

- 13 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

```

Met Ser Val Phe Val Ser Gly Ala Asn Gly Phe Ile Ala Gln His Ile
 1             5             10             15

Val Asp Leu Leu Leu Lys Glu Asp Tyr Lys Val Ile Gly Ser Ala Arg
      20             25             30

Ser Gln Glu Lys Ala Glu Asn Leu Thr Glu Ala Phe Gly Asn Asn Pro
      35             40             45

Lys Phe Ser Met Glu Val Val Pro Asp Ile Ser Lys Leu Asp Ala Phe
      50             55             60

Asp His Val Phe Gln Lys His Gly Lys Asp Ile Lys Ile Val Leu His
      65             70             75             80

Thr Ala Ser Pro Phe Cys Phe Asp Ile Thr Asp Ser Glu Arg Asp Leu
      85             90             95

Leu Ile Pro Ala Val Asn Gly Val Lys Gly Ile Leu His Ser Ile Lys
      100            105            110

Lys Tyr Ala Ala Asp Ser Val Glu Arg Val Val Leu Thr Ser Ser Tyr
      115            120            125

Ala Ala Val Phe Asp Met Ala Lys Glu Asn Asp Lys Ser Leu Thr Phe
      130            135            140

Asn Glu Glu Ser Trp Asn Pro Ala Thr Trp Glu Ser Cys Gln Ser Asp
      145            150            155            160

Pro Val Asn Ala Tyr Cys Gly Ser Lys Lys Phe Ala Glu Lys Ala Ala
      165            170            175

Trp Glu Phe Leu Glu Glu Asn Arg Asp Ser Val Lys Phe Glu Leu Thr
      180            185            190

Ala Val Asn Pro Val Tyr Val Phe Gly Pro Gln Met Phe Asp Lys Asp
      195            200            205

Val Lys Lys His Leu Asn Thr Ser Cys Glu Leu Val Asn Ser Leu Met
      210            215            220

His Leu Ser Pro Glu Asp Lys Ile Pro Glu Leu Phe Gly Gly Tyr Ile
      225            230            235            240

Asp Val Arg Asp Val Ala Lys Ala His Leu Val Ala Phe Gln Lys Arg
      245            250            255

Glu Thr Ile Gly Gln Arg Leu Ile Val Ser Glu Ala Arg Phe Thr Met

```

- 14 -

260	265	270
Gln Asp Val Leu Asp Ile Leu Asn Glu Asp Phe Pro Val Leu Lys Gly		
275	280	285
Asn Ile Pro Val Gly Lys Pro Gly Ser Gly Ala Thr His Asn Thr Leu		
290	295	300
Gly Ala Thr Leu Asp Asn Lys Lys Ser Lys Lys Leu Leu Gly Phe Lys		
305	310	315
Phe Arg Asn Leu Lys Glu Thr Ile Asp Asp Thr Ala Ser Gln Ile Leu		
325	330	335
Lys Phe Glu Gly Arg Ile		
340		

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1026 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

AUGUCAGUUU UCGUUUCAGG UGCUAACGGG UUCAUUGCCC AACACAUUGU CGAUCUCCUG	60
UUGAAGGAAG ACUAUAAGGU CAUCGGUUCU GCCAGAAGUC AAGAAAAGGC CGAGAAUUUA	120
ACGGAGGCCU UUGGUAACAA CCCAAAUUC UCCAUGGAAG UUGUCCAGA CAUAUCUAAG	180
CUGGACGCAU UUGACCAUGU UUUCCAAAAG CACGGCAAGG AUAUCAAGAU AGUUCUACAU	240
ACGGCCUCUC CAUUCUGCUU UGAUAUCACU GACAGUGAAC GCGAUUUUUU AADUCCUGCU	300
GUGAACGGUG UUAAGGGAAU UCUCACUCA AUUAAAAAAU ACGCCGCUGA UUCUGUAGAA	360
CGUGUAGUUC UCACCUUCUUC UUAUGCAGCU GUGUUCGAUA UGGCAAAGA AAACGAUAAG	420
UCUUUAACAU UUAACGAAGA AUCCUGGAAC CCAGCUACCU GGGAGAGUUG CCAAAGUGAC	480
CCAGUUAACG CCUACUGUGG UUCUAAGAAG UUUGCUGAAA AAGCAGCUUG GGAAUUUCUA	540
GAGGAGAAUA GAGACUCUGU AAAAUUCGAA UUAACUGCCG UUAACCCAGU UUACGUUUUU	600

- 15 -

```

GGUCCGCAAA UGUUUGACAA AGAUGUGAAA AAACACUUGA ACACAUCUUG CGAACUCGUC      660
AACAGCUUGA UGCAUUUAUC ACCAGAGGAC AAGAUACCGG AACUAUUUGG UGGAUACAUU      720
GAUGUUCGUG AUGUUGCAAA GGCUCAUUUA GUUGCCUUCU AAAAGAGGGA AACAAUUGGU      780
CAAAGACUAA UCGUAUCGGA GGCCAGAUUU ACUAUGCAGG AUGUUCUCGA UAUCCUUAAC      840
GAAGACUUCU CUGUUCUAAA AGGCAAUAUU CCAGUGGGGA AACCAGGUUC UGGUGCUACC      900
CAUAACACCC UUGGUGCUAC UCUUGAUAAU AAAAAGAGUA AGAAAUUGUU AGGUUUC AAG      960
UUCAGGAACU UGAAAGAGAC CAUUGACGAC ACUGCCUCCC AAUUUUAAA AUUUGAGGGC     1020
AGAAUA                                           1026

```

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1041 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS
- (B) LOCATION: 1..1041
- (D) OTHER INFORMATION: *S. cerevisiae* YGL157W

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

```

ATG ACT ACT GAT ACC ACT GTT TTC GTT TCT GGC GCA ACC GGT TTC ATT      48
Met Thr Thr Asp Thr Thr Val Phe Val Ser Gly Ala Thr Gly Phe Ile
  1             5             10             15

GCT CTA CAC ATT ATG AAC GAT CTG TTG AAA GCT GGC TAT ACA GTC ATC      96
Ala Leu His Ile Met Asn Asp Leu Leu Lys Ala Gly Tyr Thr Val Ile
      20             25             30

GGC TCA GGT AGA TCT CAA GAA AAA AAT GAT GGC TTG CTC AAA AAA TTT     144
Gly Ser Gly Arg Ser Gln Glu Lys Asn Asp Gly Leu Leu Lys Lys Phe
      35             40             45

AAT AAC AAT CCC AAA CTA TCG ATG GAA ATT GTG GAA GAT ATT GCT GCT     192
Asn Asn Asn Pro Lys Leu Ser Met Glu Ile Val Glu Asp Ile Ala Ala
      50             55             60

```


- 16 -

CCA AAC GCC TTT GAT GAA GTT TTC AAA AAA CAT GGT AAG GAA ATT AAG	240
Pro Asn Ala Phe Asp Glu Val Phe Lys Lys His Gly Lys Glu Ile Lys	
65 70 75 80	
ATT GTG CTA CAC ACT GCC TCC CCA TTC CAT TTT GAA ACT ACC AAT TTT	288
Ile Val Leu His Thr Ala Ser Pro Phe His Phe Glu Thr Thr Asn Phe	
85 90 95	
GAA AAG GAT TTA CTA ACC CCT GCA GTG AAC GGT ACA AAA TCT ATC TTG	336
Glu Lys Asp Leu Leu Thr Pro Ala Val Asn Gly Thr Lys Ser Ile Leu	
100 105 110	
GAA GCG ATT AAA AAA TAT GCT GCA GAC ACT GTT GAA AAA GTT ATT GTT	384
Glu Ala Ile Lys Lys Tyr Ala Ala Asp Thr Val Glu Lys Val Ile Val	
115 120 125	
ACT TCG TCT ACT GCT GCT CTG GTG ACA CCT ACA GAC ATG AAC AAA GGA	432
Thr Ser Ser Thr Ala Ala Leu Val Thr Pro Thr Asp Met Asn Lys Gly	
130 135 140	
GAT TTG GTG ATC ACG GAG GAG AGT TGG AAT AAG GAT ACA TGG GAC AGT	480
Asp Leu Val Ile Thr Glu Glu Ser Trp Asn Lys Asp Thr Trp Asp Ser	
145 150 155 160	
TGT CAA GCC AAC GCC GTT GCC GCA TAT TGT GGC TCG AAA AAG TTT GCT	528
Cys Gln Ala Asn Ala Val Ala Ala Tyr Cys Gly Ser Lys Lys Phe Ala	
165 170 175	
GAA AAA ACT GCT TGG GAA TTT CTT AAA GAA AAC AAG TCT AGT GTC AAA	576
Glu Lys Thr Ala Trp Glu Phe Leu Lys Glu Asn Lys Ser Ser Val Lys	
180 185 190	
TTC ACA CTA TCC ACT ATC AAT CCG GGA TTC GTT TTT GGT CCT CAA ATG	624
Phe Thr Leu Ser Thr Ile Asn Pro Gly Phe Val Phe Gly Pro Gln Met	
195 200 205	
TTT GCA GAT TCG CTA AAA CAT GGC ATA AAT ACC TCC TCA GGG ATC GTA	672
Phe Ala Asp Ser Leu Lys His Gly Ile Asn Thr Ser Ser Gly Ile Val	
210 215 220	
TCT GAG TTA ATT CAT TCC AAG GTA GGT GGA GAA TTT TAT AAT TAC TGT	720
Ser Glu Leu Ile His Ser Lys Val Gly Gly Glu Phe Tyr Asn Tyr Cys	
225 230 235 240	
GGC CCA TTT ATT GAC GTG CGT GAC GTT TCT AAA GCC CAC CTA GTT GCA	768
Gly Pro Phe Ile Asp Val Arg Asp Val Ser Lys Ala His Leu Val Ala	
245 250 255	
ATT GAA AAA CCA GAA TGT ACC GGC CAA AGA TTA GTA TTG AGT GAA GGT	816
Ile Glu Lys Pro Glu Cys Thr Gly Gln Arg Leu Val Leu Ser Glu Gly	
260 265 270	
TTA TTC TGC TGT CAA GAA ATC GTT GAC ATC TTG AAC GAG GAA TTC CCT	864
Leu Phe Cys Cys Gln Glu Ile Val Asp Ile Leu Asn Glu Glu Phe Pro	
275 280 285	

- 17 -

CAA TTA AAG GGC AAG ATA GCT ACA GGT GAA CCT GCG ACC GGT CCA AGC	912
Gln Leu Lys Gly Lys Ile Ala Thr Gly Glu Pro Ala Thr Gly Pro Ser	
290 295 300	
TTT TTA GAA AAA AAC TCT TGC AAG TTT GAC AAT TCT AAG ACA AAA AAA	960
Phe Leu Glu Lys Asn Ser Cys Lys Phe Asp Asn Ser Lys Thr Lys Lys	
305 310 315 320	
CTA CTG GGA TTC CAG TTT TAC AAT TTA AAG GAT TGC ATA GTT GAC ACC	1008
Leu Leu Gly Phe Gln Phe Tyr Asn Leu Lys Asp Cys Ile Val Asp Thr	
325 330 335	
GCG GCG CAA ATG TTA GAA GTT CAA AAT GAA GCC	1041
Ala Ala Gln Met Leu Glu Val Gln Asn Glu Ala	
340 345	

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 347 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met Thr Thr Asp Thr Thr Val Phe Val Ser Gly Ala Thr Gly Phe Ile	
1 5 10 15	
Ala Leu His Ile Met Asn Asp Leu Leu Lys Ala Gly Tyr Thr Val Ile	
20 25 30	
Gly Ser Gly Arg Ser Gln Glu Lys Asn Asp Gly Leu Leu Lys Lys Phe	
35 40 45	
Asn Asn Asn Pro Lys Leu Ser Met Glu Ile Val Glu Asp Ile Ala Ala	
50 55 60	
Pro Asn Ala Phe Asp Glu Val Phe Lys Lys His Gly Lys Glu Ile Lys	
65 70 75 80	
Ile Val Leu His Thr Ala Ser Pro Phe His Phe Glu Thr Thr Asn Phe	
85 90 95	
Glu Lys Asp Leu Leu Thr Pro Ala Val Asn Gly Thr Lys Ser Ile Leu	
100 105 110	
Glu Ala Ile Lys Lys Tyr Ala Ala Asp Thr Val Glu Lys Val Ile Val	
115 120 125	
Thr Ser Ser Thr Ala Ala Leu Val Thr Pro Thr Asp Met Asn Lys Gly	
130 135 140	

- 18 -

Asp Leu Val Ile Thr Glu Glu Ser Trp Asn Lys Asp Thr Trp Asp Ser
 145 150 155 160
 Cys Gln Ala Asn Ala Val Ala Ala Tyr Cys Gly Ser Lys Lys Phe Ala
 165 170 175
 Glu Lys Thr Ala Trp Glu Phe Leu Lys Glu Asn Lys Ser Ser Val Lys
 180 185 190
 Phe Thr Leu Ser Thr Ile Asn Pro Gly Phe Val Phe Gly Pro Gln Met
 195 200 205
 Phe Ala Asp Ser Leu Lys His Gly Ile Asn Thr Ser Ser Gly Ile Val
 210 215 220
 Ser Glu Leu Ile His Ser Lys Val Gly Gly Glu Phe Tyr Asn Tyr Cys
 225 230 235 240
 Gly Pro Phe Ile Asp Val Arg Asp Val Ser Lys Ala His Leu Val Ala
 245 250 255
 Ile Glu Lys Pro Glu Cys Thr Gly Gln Arg Leu Val Leu Ser Glu Gly
 260 265 270
 Leu Phe Cys Cys Gln Glu Ile Val Asp Ile Leu Asn Glu Glu Phe Pro
 275 280 285
 Gln Leu Lys Gly Lys Ile Ala Thr Gly Glu Pro Ala Thr Gly Pro Ser
 290 295 300
 Phe Leu Glu Lys Asn Ser Cys Lys Phe Asp Asn Ser Lys Thr Lys Lys
 305 310 315 320
 Leu Leu Gly Phe Gln Phe Tyr Asn Leu Lys Asp Cys Ile Val Asp Thr
 325 330 335
 Ala Ala Gln Met Leu Glu Val Gln Asn Glu Ala
 340 345

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1041 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- 19 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

AUGACUACUG AUACCACUGU UUUCGUUUCU GCGCAACCG GUUUCAUUGC UCUACACAUU	60
AUGAACGAUC UGUUGAAAGC UGGCUAUACA GUCAUCGGCU CAGGUAGAUC UCAAGAAAAA	120
AAUGAUGGCU UGCUCAAAAA AUUUAUAAC AAUCCCAAAC UAUCGAUGGA AAUUGUGGAA	180
GAUUAUUGCUG CUCCAAACGC CUUUGAUGAA GUUUUCAAAA AACAUGGUAA GGAAAUUAAG	240
AUUGUGCUAC ACACUGCCUC CCCAUUCCAU UUUGAAACUA CCAAUUUUGA AAAGGAUUUA	300
CUAACCCUG CAGUGAACGG UACAAAUCU AUCUUGGAAG CGAUUAAAAA AUAUGCUGCA	360
GACACUGUUG AAAAAGUUU UGUUACUUCG UCUACTUGCUG CUCUGGUGAC ACCUACAGAC	420
AUGAACAAAG GAGAUUUGGU GAUCACGGAG GAGAGUUGGA AUAAGGAUAC AUGGGACAGU	480
UGUCAAGCCA ACGCCGUUGC CGCAUAUUGU GGCUCGAAAA AGUUUGCUGA AAAAACUGCU	540
UGGGAAUUUC UUAAAGAAAA CAAGUCUAGU GUCAAAUUC ACUAUCCAC UAUCAAUCCG	600
GGAUUCGUUU UUGGUCCUCA AAUGUUUGCA GAUUCGCUAA AACAUGGCAU AAUACCUCC	660
UCAGGGAUCG UAUCUGAGUU AAUUCAUUCC AAGGUAGGUG GAGAAUUUUA UAAUUACUGU	720
GGCCCAUUUA UUGACGUGCG UGACGUUUCU AAAGCCCACC UAGUUGCAAU UGAAAAACCA	780
GAAUGUACCG GCCAAAGAUU AGUAUUGAGU GAAGGUUUUAU UCUGCUGUCA AGAAAUCGUU	840
GACAUCUGA ACGAGGAAU CCCUCAAUUA AAGGGCAAGA UAGCUACAGG UGAACCUCCG	900
ACCGGUCCAA GCUUUUAGA AAAAAACUCU UGCAAGUUUG ACAAUUCUAA GACAAAAAAA	960
CUACUGGGAU UCCAGUUUUA CAAUUUAAAG GAUUGCAUAG UUGACACCGC GCGCAAAUG	1020
UUAGAAGUUC AAAAUGAAGC C	1041

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1044 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- 20 -

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 1..1044

(D) OTHER INFORMATION: *S. cerevisiae* YGL039W

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

ATG ACT ACT GAA AAA ACC GTT GTT TTT GTT TCT GGT GCT ACT GGT TTC	48
Met Thr Thr Glu Lys Thr Val Val Phe Val Ser Gly Ala Thr Gly Phe	
1 5 10 15	
ATT GCT CTA CAC GTA GTG GAC GAT TTA TTA AAA ACT GGT TAC AAG GTC	96
Ile Ala Leu His Val Val Asp Asp Leu Leu Lys Thr Gly Tyr Lys Val	
20 25 30	
ATC GGT TCG GGT AGG TCC CAA GAA AAG AAT GAT GGA TTG CTG AAA AAA	144
Ile Gly Ser Gly Arg Ser Gln Glu Lys Asn Asp Gly Leu Leu Lys Lys	
35 40 45	
TTT AAG AGC AAT CCC AAC CTT TCA ATG GAG ATT GTC GAA GAC ATT GCT	192
Phe Lys Ser Asn Pro Asn Leu Ser Met Glu Ile Val Glu Asp Ile Ala	
50 55 60	
GCT CCA AAC GCT TTT GAC AAA GTT TTT CAA AAG CAC GGC AAA GAG ATC	240
Ala Pro Asn Ala Phe Asp Lys Val Phe Gln Lys His Gly Lys Glu Ile	
65 70 75 80	
AAG GTT GTC TTG CAC ATA GCT TCT CCG GTT CAC TTC AAC ACC ACT GAT	288
Lys Val Val Leu His Ile Ala Ser Pro Val His Phe Asn Thr Thr Asp	
85 90 95	
TTC GAA AAG GAT CTG CTA ATT CCT GCT GTG AAT GGT ACC AAG TCC ATT	336
Phe Glu Lys Asp Leu Leu Ile Pro Ala Val Asn Gly Thr Lys Ser Ile	
100 105 110	
CTA GAA GCA ATC AAA AAT TAT GCC GCA GAC ACA GTC GAA AAA GTC GTT	384
Leu Glu Ala Ile Lys Asn Tyr Ala Ala Asp Thr Val Glu Lys Val Val	
115 120 125	
ATT ACT TCT TCT GTT GCT GCC CTT GCA TCT CCC GGA GAT ATG AAG GAC	432
Ile Thr Ser Ser Val Ala Ala Leu Ala Ser Pro Gly Asp Met Lys Asp	
130 135 140	
ACT AGT TTC GTT GTC AAT GAG GAA AGT TGG AAC AAA GAT ACT TGG GAA	480
Thr Ser Phe Val Val Asn Glu Glu Ser Trp Asn Lys Asp Thr Trp Glu	
145 150 155 160	
AGT TGT CAA GCT AAC GCG GTT TCC GCA TAC TGT GGT TCC AAG AAA TTT	528
Ser Cys Gln Ala Asn Ala Val Ser Ala Tyr Cys Gly Ser Lys Lys Phe	
165 170 175	
GCT GAA AAA ACT GCT TGG GAT TTT CTC GAG GAA AAC CAA TCA AGC ATC	576
Ala Glu Lys Thr Ala Trp Asp Phe Leu Glu Glu Asn Gln Ser Ser Ile	
180 185 190	

- 21 -

AAA TTT ACG CTA TCA ACC ATC AAC CCA GGA TTT GTT TTT GGC CCT CAG	624
Lys Phe Thr Leu Ser Thr Ile Asn Pro Gly Phe Val Phe Gly Pro Gln	
195 200 205	
CTA TTT GCC GAC TCT CTT AGA AAT GGA ATA AAT AGC TCT TCA GCC ATT	672
Leu Phe Ala Asp Ser Leu Arg Asn Gly Ile Asn Ser Ser Ser Ala Ile	
210 215 220	
ATT GCC AAT TTG GTT AGT TAT AAA TTA GGC GAC AAT TTT TAT AAT TAC	720
Ile Ala Asn Leu Val Ser Tyr Lys Leu Gly Asp Asn Phe Tyr Asn Tyr	
225 230 235 240	
AGT GGT CCT TTT ATT GAC GTT CGC GAT GTT TCA AAA GCT CAT TTA CTT	768
Ser Gly Pro Phe Ile Asp Val Arg Asp Val Ser Lys Ala His Leu Leu	
245 250 255	
GCA TTT GAG AAA CCC GAA TGC GCT GGC CAA AGA CTA TTC TTA TGT GAA	816
Ala Phe Glu Lys Pro Glu Cys Ala Gly Gln Arg Leu Phe Leu Cys Glu	
260 265 270	
GAT ATG TTT TGC TCT CAA GAA GCG CTG GAT ATC TTG AAT GAG GAA TTT	864
Asp Met Phe Cys Ser Gln Glu Ala Leu Asp Ile Leu Asn Glu Glu Phe	
275 280 285	
CCA CAG TTA AAA GGC AAG ATA GCA ACT GGC GAA CCT GGT AGC GGC TCA	912
Pro Gln Leu Lys Gly Lys Ile Ala Thr Gly Glu Pro Gly Ser Gly Ser	
290 295 300	
ACC TTT TTG ACA AAA AAC TGC TGC AAG TGC GAC AAC CGC AAA ACC AAA	960
Thr Phe Leu Thr Lys Asn Cys Cys Lys Cys Asp Asn Arg Lys Thr Lys	
305 310 315 320	
AAT TTA TTA GGA TTC CAA TTT AAT AAG TTC AGA GAT TGC ATT GTC GAT	1008
Asn Leu Leu Gly Phe Gln Phe Asn Lys Phe Arg Asp Cys Ile Val Asp	
325 330 335	
ACT GCC TCG CAA TTA CTA GAA GTT CAA AGT AAA AGC	1044
Thr Ala Ser Gln Leu Leu Glu Val Gln Ser Lys Ser	
340 345	

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 348 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Met	Thr	Thr	Glu	Lys	Thr	Val	Val	Phe	Val	Ser	Gly	Ala	Thr	Gly	Phe
1				5					10					15	

- 22 -

Ile Ala Leu His Val Val Asp Asp Leu Leu Lys Thr Gly Tyr Lys Val
 20 25 30

Ile Gly Ser Gly Arg Ser Gln Glu Lys Asn Asp Gly Leu Leu Lys Lys
 35 40 45

Phe Lys Ser Asn Pro Asn Leu Ser Met Glu Ile Val Glu Asp Ile Ala
 50 55 60

Ala Pro Asn Ala Phe Asp Lys Val Phe Gln Lys His Gly Lys Glu Ile
 65 70 75 80

Lys Val Val Leu His Ile Ala Ser Pro Val His Phe Asn Thr Thr Asp
 85 90 95

Phe Glu Lys Asp Leu Leu Ile Pro Ala Val Asn Gly Thr Lys Ser Ile
 100 105 110

Leu Glu Ala Ile Lys Asn Tyr Ala Ala Asp Thr Val Glu Lys Val Val
 115 120 125

Ile Thr Ser Ser Val Ala Ala Leu Ala Ser Pro Gly Asp Met Lys Asp
 130 135 140

Thr Ser Phe Val Val Asn Glu Glu Ser Trp Asn Lys Asp Thr Trp Glu
 145 150 155 160

Ser Cys Gln Ala Asn Ala Val Ser Ala Tyr Cys Gly Ser Lys Lys Phe
 165 170 175

Ala Glu Lys Thr Ala Trp Asp Phe Leu Glu Glu Asn Gln Ser Ser Ile
 180 185 190

Lys Phe Thr Leu Ser Thr Ile Asn Pro Gly Phe Val Phe Gly Pro Gln
 195 200 205

Leu Phe Ala Asp Ser Leu Arg Asn Gly Ile Asn Ser Ser Ser Ala Ile
 210 215 220

Ile Ala Asn Leu Val Ser Tyr Lys Leu Gly Asp Asn Phe Tyr Asn Tyr
 225 230 235 240

Ser Gly Pro Phe Ile Asp Val Arg Asp Val Ser Lys Ala His Leu Leu
 245 250 255

Ala Phe Glu Lys Pro Glu Cys Ala Gly Gln Arg Leu Phe Leu Cys Glu
 260 265 270

Asp Met Phe Cys Ser Gln Glu Ala Leu Asp Ile Leu Asn Glu Glu Phe
 275 280 285

Pro Gln Leu Lys Gly Lys Ile Ala Thr Gly Glu Pro Gly Ser Gly Ser
 290 295 300

Thr Phe Leu Thr Lys Asn Cys Cys Lys Cys Asp Asn Arg Lys Thr Lys

- 23 -

305	310	315	320
Asn Leu Leu Gly Phe Gln Phe Asn Lys Phe Arg Asp Cys Ile Val Asp			
325	330	335	
Thr Ala Ser Gln Leu Leu Glu Val Gln Ser Lys Ser			
340	345		

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1044 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AUGACUACUG AAAAAACCGU UGUUUUGUU UCUGGUGCUA CUGGUUUC AU UGCUCUACAC	60
GUAGUGGACG AUUUAUUA AAACUGGUUAC AAGGUCAUCG GUUCGGGUAG GUCCCAAGAA	120
AAGAAUGAUG GAUUGCUGAA AAAAUUUAAG AGCAAUCCCA ACCUUUCAU GGAGAUUGUC	180
GAAGACAUUG CUGCUCCAAA CGCUUUUGAC AAAGUUUUUC AAAAGCACGG CAAAGAGAUC	240
AAGGUUGUCU UGCACAUAGC UUCUCCGGUU CACUUAACA CCACUGAUUU CGAAAAGGAU	300
CUGCUAAUUC CUGCUGUGAA UGGUACCAAG UCCAUCUAG AAGCAAUCAA AAUUAUGCC	360
GCAGACACAG UCGAAAAAGU CGUUAUUACU UCUUCUGUUG CUGCCCUUGC AUCUCCCGGA	420
GAUAUGAAGG ACACUAGUUU CGUUGUCAAU GAGGAAAGUU GGAACAAAGA UACUUGGGAA	480
AGUUGUCAAG CUAACGCGGU UUCCGCAUAC UGUGGUUCCA AGAAAUUUGC UGAAAAACU	540
GCUUGGGAUU UUCUCGAGGA AAACCAAUCA AGCAUCAAU UUACGCUAUC AACCAUCAAC	600
CCAGGAUUUG UUUUUGGCCC UCAGCUAUUU GCCGACUCUC UUAGAAAUUG AAUAAUAGC	660
UCUUCAGCCA UUAUUGCCAA UUUGGUUAGU UAUAAAUAG GCGACAAUUU UUAUAAUAC	720
AGUGGUCCUU UUAUUGACGU UCGCGAUGUU UCAAAAGCUC AUUUACUUGC AUUUGAGAAA	780
CCCGAAUGCG CUGGCCAAAG ACUAUUCUUA UGUGAAGUA UGUUUUGCUC UCAAGAAGCG	840

- 24 -

CUGGAUAUCU UGAAUGAGGA AUUCCACAG UUAAAAGGCA AGAUAGCAAC UGGCGAACCU	900
GGUAGCGGCU CAACUUUUU GACAAAAAAC UGCUGCAAGU GCGACAACCG CAAAACCAA	960
AAUUUAUUAG GAUCCAAUU UAAUAAGUUC AGAGAUUGCA UUGUCGAUAC UGCCUCGCAA	1020
UUACUAGAAG UUCAAGUAA AAGC	1044

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/23419

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12P 21/06; C12N 9/02, 1/20, 15/00; C07H 21/04; C07K 1/00.

US CL : 435/69.1, 189, 252.3, 320.1; 536/23.2, 23.7; 530/350.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.1, 189, 252.3, 320.1; 536/23.2, 23.7; 530/350.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JORNVALL et al. Short-Chain Dehydrogenases/Reductases (SDR). Biochemistry. 09 May 1995. Vol 34, No. 18, pages 6003-6013, see the entire article.	1-21

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 29 DECEMBER 1998	Date of mailing of the international search report 03 FEB 1999
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer TEKCHAND SAIDHA Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/23419

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN Files : Medline, Caplus, Wpids, Biosis, Biotechds, Scisearch. Search terms included : ketoreductase, Zygosaccharomyces, yeast in combinations as well as authors names search. APS search.